



# ReaX ZINC PLUS

4-0-0

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....	4.0%
4.0% Ammoniacal Nitrogen .....	
Sulfur (S) .....	3.0%
3.0% Combined Sulfur (S) .....	
Zinc (Zn) .....	6.5%
6.5% Water Soluble Zinc (Zn) .....	

**Derived From:** Anhydrous Ammonia, Zinc Sulfate

### ALSO CONTAINS NON-PLANT FOOD INGREDIENT

4.00% .....	Organic Matter (derived from leonardite)
96.00% .....	Total Other Ingredients (inert as a nonplant food ingredient)

**KEEP OUT OF REACH OF CHILDREN**



## WARNING

Avoid breathing mists. Harmful if swallowed. Avoid contact with skin, eyes or clothing.

### FIRST AID

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. **If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT, CALL 1-866-944-8565.**

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk or street back onto your lawn.

### DIRECTIONS FOR USE

Uses include but are not limited to the research verified crop specific directions in the table below.

CROP	APPLICATION RATE PER ACRE	APPLICATION DIRECTIONS AND TIMING
(01) Root and Tuber Vegetables including Carrot, Ginseng, Horseradish, Parsley (turnip-rooted), Potato, Radish, Sugar Beet, Sweet Potato, Turnip	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(03) Bulb Vegetables including Garlic, Leek, Onion	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(04) Leafy Vegetables including Celery, Head and Leaf Lettuce and Spinach	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(05) Brassica Vegetables including Broccoli, Cabbage, Cauliflower, Kale, and Mustard greens	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(06) Legume Vegetables (Succulent or Dried) including Bean (all types), Peas and Soybeans	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(08) Fruiting Vegetables including Eggplant, Pepper and Tomato	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(09) Cucurbit Vegetables including Cantaloupe, Cucumber, Honeydew, Muskmelon, Squash (summer and winter) and Watermelon	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(10) Citrus Fruit including Grapefruit, Lemon, Lime, Sweet Orange and Tangerine	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Soil:</b> Inject 3 to 5 inches deep at the drip line, or surface band and water into the soil. <b>Foliar:</b> Repeat applications may be beneficial. Dilute with water to ensure uniform distribution.
(11) Pome Fruits including Apple and Pear	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Soil:</b> Inject 3 to 5 inches deep at the drip line, or surface band and water into the soil. <b>Foliar:</b> Repeat applications may be beneficial. Dilute with water to ensure uniform distribution.
(12) Stone Fruits including Apricot, Cherry, Peach and Plumcot	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Soil:</b> Inject 3 to 5 inches deep at the drip line, or surface band and water into the soil. <b>Foliar:</b> Repeat applications may be beneficial. Dilute with water to ensure uniform distribution.
(13) Berry and Small Fruit including Blackberry, Blueberry, Caneberry, Kiwi, Raspberry and Grape	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Soil:</b> Inject 3 to 5 inches deep at the drip line, or surface band and water into the soil. <b>Foliar:</b> Repeat applications may be beneficial. Dilute with water to ensure uniform distribution.
(14) Tree Nuts including Almonds, Cashews and Pecans	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Soil:</b> Inject 3 to 5 inches deep at the drip line, or surface band and water into the soil. <b>Foliar:</b> Repeat applications may be beneficial. Dilute with water to ensure uniform distribution.
(15) Grain, Cereal	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(16) Forage, Fodder and Straw of Cereal Grains	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(17) Grass Forage, Fodder and Hay	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(18) Nongrass Animal Feeds including Alfalfa, Clover, Hay and Vetch	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(19) Herbs and Spices including Basil, Dill, Mustard and Sage	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition. <b>Foliar:</b> Dilute with water to ensure uniform distribution.

### Directions for use cont'd.

CROP	APPLICATION RATE PER ACRE	APPLICATION DIRECTIONS AND TIMING
(20) Oil Seed Crops including Canola, Flax and Sunflower	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
(22) Stalk, Stem and leaf petiole vegetables, Asparagus, bamboo shoots and celery	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	<b>Caution:</b> Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.
(99) Cotton, Peanut, Sugarcane, Tobacco	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
Ornamental and Landscape	<b>Soil:</b> 1.0 to 8.0 quarts <b>Foliar:</b> 1.0 to 4.0 quarts	
Turf	<b>Soil:</b> 0.75 to 6 oz per 1000 square feet <b>Foliar:</b> 0.75 to 3 oz per 1000 square feet	

### Soil Applications:

ReaX Zinc may be applied pre-plant, as a starter, side dress, fertigated, top dressed, or banded to the soil. To apply alone, dilute with water to ensure uniform distribution.

### Fertigation:

Can be applied through Center Pivot, Drip and Micro Sprinkler Irrigation systems.

All Crops: 1 to 8 quarts per acre.

Continue irrigation for a minimum of 2 hours after application to flush the lines and to maximize nutrient distribution with drip and micro sprinkler use.

Inject after the irrigation water filter.

### Transplant Water – All Crops:

1 to 4 quarts per acre not to exceed a 3% solution of ReaX Zinc to water.

Apply in a minimum of 50 gallons solution.

Mixing with water for field storage is generally not recommended. Products blended with water should be used immediately and not stored for prolonged periods.

### Aerial Application:

In aerial application use adequate water to obtain thorough coverage.

### Caution:

Always check compatibility with other products prior to use.

Do NOT combine with Calcium Fertilizers such as CAN-17.

Fertilizer solutions with a pH of 5 or less may not be compatible.

Combining with liquid fertilizer solutions is more effective if done at a blending plant with vigorous agitation in blending process.

### Caution:

Foliar applications close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.

### STORAGE AND DISPOSAL

**STORAGE:** Store in original containers only. Keep containers tightly closed when not in use. Store in a cool, dry well-ventilated area, preferably in a locked storage area away from children, feed and food products, and seed. Store at ambient temperatures. Do not contaminate water, food, or feed by storage or disposal. **DISPOSAL:** Do not reuse container. Completely empty container into application equipment. Thoroughly rinse container, adding rinse water to the spray tank.

For help with any spill, leak, fire or exposure involving this material, call day or night CHEMTREC 1-800-424-9300.

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

BEFORE BUYING OR USING THIS PRODUCT, read the Directions for Use and the following Conditions of Sale and Limitation of Warranty and Liability. By buying or using this product, the buyer or user accepts the following Conditions of Sale and Limitation of Warranty and Liability, which no employee or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary. LOVELAND PRODUCTS, INC. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, when the product is used in accordance with such Directions for Use under normal conditions of use. LOVELAND PRODUCTS, INC. MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THE BUYER'S OR USER'S EXCLUSIVE REMEDY FOR ANY INJURY, LOSS, OR DAMAGE RESULTING FROM THE HANDLING OR USE OF THIS PRODUCT SHALL BE LIMITED TO ONE OF THE FOLLOWING, AT THE ELECTION OF LOVELAND PRODUCTS, INC. OR THE SELLER: DIRECT DAMAGES NOT EXCEEDING THE PURCHASE PRICE OF THE PRODUCT OR REPLACEMENT OF THE PRODUCT. LOVELAND PRODUCTS, INC. AND THE SELLER SHALL NOT BE LIABLE TO THE BUYER OR USER OF THIS PRODUCT FOR ANY CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-lpi.com/>

Tests have not been carried out under all possible use conditions and Seller cannot and does not represent or warrant that the goods are compatible with all other chemicals or under all use conditions.

ReaX Plus Zinc is a registered trademark of Loveland Products, Inc.




### WARNING

This product can expose you to chemicals including Nickel, which are known to the State of California to cause cancer. For more information go to [www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov).

F1548

FOR AGRICULTURAL AND PROFESSIONAL USE ONLY



Density: 11.8 lbs/gallon @ 68 °F

**Net Contents: 265 gallons (1,003.13 liters)**

**Net Weight: 3,127 lbs (1,418.38 kg)**

Product Code:



Density: 11.8 lbs/gallon @ 68 °F

**Net Contents:**                      gallons (                      liters)

**Net Weight:**                      lbs (                      kg)

Product Code: